



Review Article

ANTIMICROBIAL ACTIVITY OF 1, 3, 4-OXADIAZOLES: A REVIEW


Himanshu¹ and Vivek Kumar Joshi^{2*}*Corresponding Author: Vivek Kumar Joshi ✉ misra.prem@rediffmail.com

This review provides readers with an overview of their broad spectrum of pharmacological activities for 1, 3, 4-oxadiazole derivatives, and as reported over the past years. The search for new antimicrobial agents will consequently always remain as an important and challenging task for medicinal chemists. This Review has basic information about 1, 3, 4-oxadiazole and its antimicrobial activity work for further development in this field.

Keywords: 1, 3, 4-oxadiazole, Antimicrobial activity, Review

INTRODUCTION

Oxadiazole is a heterocyclic aromatic compound of molecular formula $C_2H_2N_2O$. It is a five membered ring consisting of 2 nitrogen atoms, 2 carbon atoms, 1 oxygen atom and 2 double bonds which shows antimicrobial, anticancer, anti-inflammatory and antioxidant activities etc. . This review article has summarized vital information on antimicrobial activity of 1, 3, 4-oxadiazole heterocyclic nucleus to provide effective antimicrobial drugs by solving the problem of microbial resistance towards currently used antibiotics. 1, 3, 4-Oxadiazole is a well known heterocyclic nucleus because of its broad spectrum of pharmacological activities especially potent antimicrobial activities. There are 4 isomers of oxadiazole as shown in the Figures below.



Oxadiazole shows inductive effect because of the presence of heteroatom in the ring and thus it is considered to be a weak base. It consists of 2 pyridine like nitrogen, due to which it exhibits conjugate diene type character. Electrophilic substitution at carbon is very difficult in this case due to less electron density which is mainly due to the presence of pyridine like nitrogen in the ring that shows electron withdrawal effect.

Due to the presence of two pyridine type nitrogen, the aromaticity will be removed. Many studies on comparison between 1, 2, 4- and 1, 3, 4-oxadiazole pairs shows that, in all cases, 1, 3, 4-oxadiazole isomer shows lower magnitude lipophilicity as compared to its isomeric partner.

¹ Meerut Institute of Engineering and Technology, Meerut, Uttar Pradesh 250005, India.

² Mewar University, Rajasthan, India.

Other differences involve metabolic stability, hERG inhibition, and aqueous solubility. All these studies favoured the 1, 3, 4-oxadiazole isomers. The difference in profile between the 1, 2, 4 and 1, 3, 4 regioisomers can be rationalized by their intrinsically different charge distributions. The 1, 3, 4-oxadiazole undergoes number of reactions including electrophilic substitution, nucleophilic substitution, thermal and photochemical.

2, 5-disubstituted-1,3,4-oxadiazole derivatives are colorless substances. Replacement of an alkyl residue by an aryl radical considerably raises the melting and boiling points. Usually the asymmetrical 1,3,4-oxadiazole derivatives melt and boil at lower temperature than the symmetrical compounds.

The solubility of oxadiazoles in water varies with the substituents present: 2, 5-dimethyl-1, 3, 4-oxadiazole is miscible with water in all proportions whereas the solubility of 2, 5-diphenyl-1, 3, 4-oxadiazole in water is less.

Electrophilic introduction of functional groups (for example nitro or sulphuric acid groups) into the nucleus is unusual. Electrophilic substitution occurs in aryl substituent. Halogenations is also difficult, but 2, 5-diaryl-1, 3, 4-oxadiazoles, afford complexes with halogens. A range of acylation and alkylation reactions of hydroxyl, thio and amino-1, 3, 4-oxadiazoles occur at the ring nitrogen.


Nucleophilic substitution reactions are uncommon. Many of nucleophilic reagents cause ring cleavage, depending upon the electron density at C-2 and C-5. Generally aryl substituted 1, 3, 4-oxadiazoles are less sensitive to acid than the alkyl substituted derivatives.

An antimicrobial is an agent that kills microorganisms or inhibits their growth.

Antimicrobial medicines can be grouped according to the microorganisms they act primarily against. For example, antibiotics are used against bacteria and antifungal are used against fungi. They can also be classified according to their function. Agents that kill microbes are called microbicidal, while those that merely inhibit their growth are called biostatic. The use of antimicrobial medicines to treat infection is known as antimicrobial chemotherapy, while the use of antimicrobial medicines to prevent infection is known as antimicrobial prophylaxis.


The most recent literature survey on antimicrobial potential of novel 1,3,4-oxadiazole derivatives has been presented in this section as given below:

Arvind K. Singh *et al.* Synthesis of (ethyl 2-(1H Benzo [d] [1, 2, 3] triazole –1 – yl) acetate) and (2H – benzo [d] [1, 2, 3] triazole – 1 – yl acetate to hydrazine) along with their derivatives has been done. The Antimicrobial activity of the synthesized compounds was evaluated, on *Streptococcus aureus* and *Escherichia coli*. The present investigation deals with the synthesized compounds possessing good antimicrobial activity.




Shridhar *et al* synthesized 1,3,4-oxadiazole incorporated azo dye derivatives (2) and evaluated their antimicrobial activity. The antimicrobial activity of newly synthesized compounds was determined by well plate method in nutrient agar (antibacterial activity) and

Sabouraud dextrose agar (antifungal activity). *E. coli*, *Staphylococcus aureus*, *Bacillus subtilis*, *Salmonella typhi*, were used to investigate the antibacterial activities and *Pseudomonas Aeruginosa*, *Candida albicans*, *Candida parapsilosis*, were used to investigate the antifungal activities. One compound showed maximum inhibitory activity against *Pseudomonas aeruginosa* and *Candida parapsilosis* at MIC 2.5 mg/ml. Antibacterial drug Ampicillin and antifungal drug Fluconazole were used as standard drug.




Parikh *et al.* synthesized a novel series of 4-((5-(2-hydroxyphenyl)-1,3,4-oxadiazol-2-yl)sulfonyl)benzaldehyde derivatives and screened for antimicrobial activity by filter paper disc method. Stains used were *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Bacillus subtilis*, *Escherichia coli* for the antibacterial activity and for the antifungal activity fungi *Candida albicans*. Ciprofloxacin and Ampicillin as standard antibacterial drug and Ketoconazole and Fluconazole as standard antifungal drug were used. From the result it was found that some compounds showed excellent antibacterial activity against *P. aeruginosa* whereas other


compounds displayed better antibacterial activity against *B. subtilis*.




Jun-Shu *et al.* synthesized the series of several new 5-[4'-(5-phenyl-1,3,4-oxadiazole-2-yl)sulfonylmethyl]-biphenyl-2-yl]-tetrazole derivatives and these compounds were evaluated for their antimicrobial activity against *B. subtilis* and *E. coli* at the concentration of 100 µg/ml in nutrient agar media. These compounds demonstrated significant antimicrobial activities as compared with standard drug.



Fuloria *et al.* synthesized a new series 1-(2-aryl-5-phenethyl-1,3,4-oxadiazole-3(2H)-yl)-ethanones and investigated for their antimicrobial activities. These newly synthesized compounds showed good antibacterial activity against the stains of micro-organisms like *S. aureus*, *P. aeruginosa* as compared with standard drug.




Somani *et al.* synthesized A novel series of 2-(3-chlorobenzylidene)-1-(1-(5-(pyridine-4-yl)-1,3,4-oxadiazol-2-ylthio) prop-2-en-2 yl)hydrazine derivatives and evaluated for their antimicrobial activity. The *in vitro* antibacterial activity against *S. aureus* and *E. coli* was determined by cup-plate method by using Ampicillin as standard drug. The *in vitro* antifungal activity of titled compounds was carried out against *C. albicans* and *A. niger* by using Fluconazole as standard drug. The tests were repeated thrice to confirm the findings that the some compounds exhibited good antibacterial activities against *S. aureus* while some compounds were effective against *E. coli*. One compound was found to be the most potent against *C. albicans* and two compounds were more active as antifungal agents against *A. niger* on comparison with standard drug.




Patel *et al.* synthesized 4-thiazolidinone incorporated 1, 3, 4-oxadiazoles and screened them for their antibacterial and antifungal *in vitro* activity against *E. coli*, *P.aeruginosa*, *S. aureus*, *S. pyogenes*, *C.albicans*, *A. niger*, *A. clavatus* respectively by using broth microdilution method.

Minimum inhibitory concentration was determined and compared with standard drugs Ampicillin and Griseofulvin. Some compounds showed good antimicrobial activity and other compounds depicted moderate activity.




Demirbas *et al.* synthesized 2-(5-mercapto)-1, 3, 4-oxadiazol-2-ylmethyl-1, 2, 4-triazol-3-one derivatives. The antimicrobial effects of the substances were tested quantitatively in their respective broth media by using double dilution and the Minimal Inhibition Concentration (MIC) values were determined. Ampicillin and fluconazole were used as standard antibacterial and antifungal drugs, respectively. Few compounds displayed good to moderate antimicrobial activity .




Kanthiah *et al.* synthesized (2-aminophenyl)-1, 3, 4-oxadiazole-2(3H)-thione derivatives and screened for their antimicrobial activities. Gram positive microorganisms such as *Staphylococcus aureus*, *Streptococcus pyogenes*, Gram negative microorganisms such as *Escherchia coli*,


Klebsiella aerogenes and fungus *Candida albicans* were used for antibacterial studies. The disc diffusion method was used to evaluate antimicrobial activities. Compounds at a concentration of 100 µg/ml showed good antibacterial and antifungal activities against all the tested microorganisms. Amikacin and Ketoconazole were used as standard drugs for antibacterial and antifungal activities respectively.




Farshori *et al.* synthesized 5-alkenyl/hydroxyalkenyl-2-phenylamine-1,3,4-Oxadiazoles and tested for their *in vitro* antimicrobial activities by disc diffusion method. Among the synthesized compounds, some compounds were found to be active against fungal stain i.e. *Penicillium marneffei* and was compared with Griseofulvin as standard drug whereas other compounds were found to be active against bacterial stains like *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Streptococcus pyogenes*, *Klebsiella pneumoniae* and were compared with Chloramphenicol as standard drug.




Reddy *et al.* synthesized 4-fluoroanilinomethyl, 4-chloroanilinomethyl, 2-trifluoromethylanilinomethyl-oxadiazole derivatives. The synthesized compounds were evaluated for their antibacterial activity against three representative Gram positive bacteria *Bacillus subtilis*, *Bacillus sphaericus*, *Staphylococcus aureus* and three Gram negative bacteria *Pseudomonas aeruginosa*, *Klebsiella aerogenes* and *Chromobacterium violaceum*. In addition, these compounds were also screened for their antifungal activity against four fungal microorganisms *Candida albicans*, *Aspergillus fumigatus*, *Trichophyton rubrum* and *Trichophyton mentagrophytes*. Most of the compounds showed excellent antimicrobial activities when compared with their respective standard drugs.




Mishra *et al.* synthesized A series of 3,4-dihydro-6-methyl-4-(4-nitrophenyl)-5-(5-phenyl-1,3,4-oxadiazol-2-yl) pyrimidin-2-(1H)-one derivatives and tested for their antimicrobial activity by cup and plate method. Some compounds showed promising antibacterial activity against gram positive bacteria *Streptococcus pneumoniae* and other compounds displayed promising antibacterial activity against gram positive bacteria *Escherichia coli* as compared to standard drugs Ofloxacin and Levofloxacin.




Chandrakantha *et al.* synthesized some novel 1, 3, 4-oxadiazole derivatives bearing 2-fluoro-4-methoxy phenyl moiety and screened for antimicrobial activity through serial dilution method. Amongst the various compounds synthesized, two compounds showed excellent antibacterial activity against *Escherichia coli* and *Pseudomonas aeruginosa* and other compounds displayed excellent antifungal activity against *Candida albicans*. Compounds tested for antibacterial activity were compared with standard drug Furacin and for antifungal activity standard drug was Flucanazole.




Chen *et al.* synthesized 5-(3, 4, 5-trimethoxyphenyl)-2-sulfonyl-1, 3, 4-oxadiazole derivatives and investigated for their antifungal activity against *Gibberellae*, *Botrytis cinerea*, *Sclerotinia sclerotiorum*. Amongst the tested compounds, two compounds depicted promising antifungal activities when compared with standard drug Hymexazol.




Karthikeyan *et al.* synthesized 2, 4-dichloro-5-fluorophenyl containing oxadiazoles and then final compounds were demonstrated for their antimicrobial activity. Few compounds displayed good inhibition against *Staphylococcus aureus*, *Escherichia coli* when compared with standard drug Ciprofloxacin. Some other compounds also showed good inhibition against all the fungal stains *Candida albicans*, *Aspergillus fumigatus* and *Penicillium marneffeii* when compared with standard drug Greseofluvin.




Rai *et al.* synthesized a novel series of 2-[1-(5-chloro-2-methoxyphenyl)-5-methyl-1H-pyrazol-4-yl]-5-(substitutedphenyl)-1, 3, 4-oxadiazole derivatives and investigated for their antibacterial activity. From the tested compounds, few compounds showed significant activity against *Bacillus subtilis* and *Escherichia coli*, *Staphylococcus aureus* and *Klebsiella pneumonia*. These compounds were compared with Ampicillin as standard drug.




Li *et al.* synthesized (E)- α -(methoxyimino)-benzeneacetate derivatives containing 1, 3, 4-Oxadiazole ring and investigated for their fungicidal activities. All of the compounds exhibited significant fungicidal activities against *Rhizoctonia solani*, *Botrytis cinereapers*, *Gibberealpers zaeae*, *Physalospora piricola* and *Bipolaris mayclis* when compared with standard drug.




Bhardwaj *et al.* synthesized a novel series of 2-(5-(3-chlorophenyl)-1,3,4-oxadiazol-2-yl)-1H-indole derivatives and demonstrated for their antimicrobial activity on different stains. Out of total of compounds synthesized, three compounds were found to be active against bacterial stains like *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas aeruginosa* whereas none of the compound was found to be effective against fungal stains. Standard Drugs used for comparison of antimicrobial activities were Norfloxacin and Fluconazole.




Prakash *et al.* synthesized a series of novel unsymmetrical 2-[1, 3-bis(4-nitrophenyl)-1H-pyrazol-4-yl]-5-phenyl-1, 3, 4-oxadiazoles and tested for their antibacterial and antifungal activities. Amongst the tested compounds, two compounds depicted most potent antibacterial activity against *Staphylococcus aureus* and was compared with ciprofloxacin as standard drug. Two compounds displayed maximum inhibition against both of the fungi *Aspergillus niger* and *Aspergillus flavus* when compared with Fluconazole as standard drug.




Mayekar *et al.* synthesized a series of new 5-amino, 2-(5-bromo) naphthalene) carboxalate, 1, 3, 4-oxadiazole derivatives having 6-bromonaphthalene moiety. The antimicrobial activities of title compounds were examined against two gram positive bacteria (*S. aureus*, *S. pyogenes*), two gram negative bacteria (*E. coli*, *P. aeruginosa*) and three fungi (*C. albicans*, *A. niger*, *A. clavatus*) by using the broth microdilution method. Some of the tested compounds displayed significant antimicrobial activities when compared with their respective standard drugs.



Ansari *et al.* synthesized some new 3-acetyl-5-(3-chloro-1-benzo[b]thiophen-2-yl)-2-substituted phenyl-2, 3- dihydro-1, 3, 4-oxadiazoles (22) and evaluated for their antimicrobial activities. All the compounds were screened for their antibacterial activities against *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli* and *Pseudomonas aeruginosa* and for antifungal activity against *Candida albicans* and *Asperigillus niger*. Some compounds exhibited significant antibacterial and moderate antifungal activities on comparison with their respective standard drugs.




Dewangan *et al.* synthesized novel derivatives of 4-(5-chloro-1, 3, 4-oxadiazole-2-yl) pyridine and evaluated for antimicrobial activities. The antimicrobial activities of title compounds were examined against two gram positive bacteria (*S. aureus*, *S. pyogenes*), two gram negative bacteria (*E. coli*, *P. aeruginosa*) and three fungi (*C. albicans*, *A. niger*, *A. clavatus*) using the broth microdilution method. Some derivatives exhibited excellent antimicrobial activities on comparison with their respective standard drugs.




Prakash *et al.* synthesized a novel series of 5-(amino(4-bromophenyl)methyl)-3-((4-phenylpiperazin-1-yl)methyl)-1, 3, 4-oxadiazole-


2(3H) thiones. The newly synthesized compounds were investigated for their antimicrobial activities. Microbial stains used for antibacterial activity were *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichiacoli* and *Pseudomonas aeruginosa* and for antifungal activity *Candida albicans* and *Asperigillus niger*. Few compounds showed good antimicrobial activity on comparison with standard drug Norfloxin.




Kakanejadifard *et al.* synthesized some novel 2-((4-amino-1, 2, 5-oxadiazol-3- ylimino)methyl)-4-(phenyldiazenyl) phenol derivatives and evaluated their antimicrobial activities. The synthesized compounds were screened for their *in vitro* antimicrobial activity against both Gram-positive (*Staphylococcus aureus* and *Bacillus cereuss*) and Gram negative (*Escherichia coli* and *Klebsiella pneumonia*) bacteria by using disc diffusion method. The antibacterial activity was reported as the minimum inhibitory concentration (MIC) in mg/ml. One compound showed the most potent antimicrobial activity with MIC value of 57 mg/ml against *S. aureus* and *B. cereuss* as compared with standard drug.




Kadi *et al.* synthesized a novel series of 2-(1-adamantylamino)-5-substituted-1, 3, 4-oxadiazole derivatives and screened their antimicrobial activity. The primary screening was carried out using the agar disc diffusion method using Muller Hinton agar medium. Antimicrobial activity of compounds was compared with antibacterial drug Ampicillin and the antifungal drug Clotrimazole against *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans*. Some compounds depicted good to moderate antimicrobial activity relative to standard drug.



Shaker *et al.* synthesized N-phenyl-5-(4-(5-phenylamino-1, 3, 4-oxadiazol-2-yl) phenyl)-1, 3, 4-oxadiazol-2-amine and examined their antimicrobial activity. Gram-negative bacteria (*Serratia*), gram-positive bacteria (*Bacillus cereus*), as well as two different fungi, *Fusarium moniliforme* and *Aspergillus flavus* were used for this purpose. Some compounds showed good antimicrobial activity whereas other compounds displayed moderate antimicrobial activity as compared with standard drug.




Rahman *et al.* synthesized compounds some new (3a,7a-dihydro-1H-indol-3-yl) (4-methyl- 1, 3, 4-oxadiazolidin-3-yl) methanone derivatives and screened *in vitro* for their antimicrobial activities against four stains of bacteria (*Staphylococcus aureus*, *Serratia marcescens*, *Streptococcus*, *Pseudomonas aeruginosa*) and two species of fungi (*Aspergillus parasiticus*, *Penicillium oxalicum*) using the filter paper disc method. Most of the compounds exhibited considerable activities against two bacterial species, *Serratia marcescens* and *Streptococcus*. Some compounds exhibited a moderate activity against *Staphylococcus aureus*. All the screened compounds were inactive against *Pseudomonas aeruginosa*. Some compounds showed moderate antibacterial activity against *Penicillium oxalicum*.



Nazk Mohammed Aziz *et al.* synthesized new metal complexes of the ligand (HL) 2-[1H-Pyrrol-2-ylimino methyl]-5-phenyl-1, 3, 4-oxadiazol with the metal ions Co(II), Ni(II) and Cu(II), were prepared in alcoholic medium. The Schiff bases were condensed by using [Pyrrolcarboxaldehyde] with [2-amino-5-(phenyl-1, 3, 4-oxadiazole)] in alcoholic medium. As the Schiff base prepared was tridentate ligand, it was used for forming complexes with Co+2, Ni+2, Cu +2 and Zn+2 ions of type M (HL)₂. All the synthesized Schiff base and their metal complexes were characterized

by FTIR Spectroscopy, Electronic Spectroscopy, Elemental Analysis, Magnetic Susceptibility Measurements, Thermal Analysis, ^1H -NMR Spectra, and Mass Spectra. The antimicrobial activity of these compounds was determined by the agar diffusion method. These types of bacteria *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans* were used to show the biological activities of the ligand and its complexes.



CONCLUSION

Various oxadiazole derivatives have been placed a successfully used antimicrobial drugs in clinical practice. This review has highlighted the use of Oxadiazole derivatives having antimicrobial activity. The 1, 3, 4-oxadiazole have shown significant antimicrobial activity against a wide range of microorganisms. In conclusion we arrive at the postulate that 1, 3, 4-oxadiazole has antimicrobial activity. An important antimicrobial drug is Furazolidone.

Furazolidone is a nitrofurantoin antibacterial agent. It is marketed by Roberts Laboratories under the brand name Furoxone and by GlaxoSmithKline as Depandal-M, Diafuron, medaron. Furazolidone as an antimicrobial is used to treat diarrhea or enteritis caused by bacteria and protozoa including traveler's diarrhea, food poisoning, typhoid fever, cholera, salmonella infections, and giardiasis.

ACKNOWLEDGMENT

The first author is specially thankful to Mr. Prem

Shankar (Asst. Professor MIET Baghpat Bypass Meerut) for the guidance and Support and for providing all the possible facilities for the research work.

REFERENCES

1. Afsharin Zarghi, Samaneh Hamed, Fatemeh Tootooni, Behzad Amini, Behrang Sharifi, Mehrdad Faizi, Seyed Abbas Tabatabai and Abbas Shafiee (2008), "Synthesis and Pharmacological Evaluation of New 2-Substituted-5-{2-[(2-halobenzyl) thio] phenyl}-1, 3, 4-oxadiazoles as Anticonvulsant Agents", *Sci Pharm.*, Vol. 76, pp. 185-201.
2. Anil N Mayekar, Yathirajan H S, Narayana B, Sarojini B K and Suchetha Kumari N (2010), "Synthesis and Antimicrobial Studies on New Substituted 1, 3, 4-Oxadiazole Derivatives Bearing 6-Bromonaphthalene Moiety", *International Journal of Chemistry*, Vol. 2, No. 2.
3. Ansari K F and Lal C (2009), "Synthesis, Physicochemical Properties and Antimicrobial Activity of Some New Benzimidazole Derivatives", *Eur J Med Chem*, Vol. 44, pp. 4028-4033.
4. Ansari K F and Lal C (2009), "Synthesis, Physicochemical Properties and Antimicrobial Activity of Some New Benzimidazole Derivatives", *European Journal of Medicinal Chemistry*, Vol. 44, pp. 4028-4033.
5. Archer J S and Archer D F (2002), "Oral Contraceptive Efficacy and Antibiotic Interaction: A Myth Debunked", *Journal of the American Academy of Dermatology*, Vol. 46, No. 6, pp. 917-923.

6. Asif Hussain and Mohammed Ajmal (2009), "Synthesis of Novel 1, 3, 4-oxadiazoles Derivatives and their Biological Properties", *Acta Pharm.*, Vol. 59, pp. 223-233.
7. Ates O, Kocabalkanli A, Cesur N and Otuk G (1998), "Synthesis and Antimicrobial Activity of Some 5-aryl-2-[N, N-disubstituted thiocarbamoylthio) Acylamino]-1, 3, 4-oxadiazoles", *Farmaco II*, Vol. 53, No. 8, pp. 541-544.
8. Aziz-ur-Rehman, Asia Siddiqua, Muhammad AAbbasi, Shahid Rasool, Sabahat Z Siddiqui, Irshad Ahmad and Saira Afzal (2014), "Synthesis of Some New 5-substituted-2-((6-chloro-3, 4 methylenedioxyphenyl) methylthio)-1, 3, 4- Oxadiazole Derivatives as Suitable Antibacterial Inhibitors", *Bulletin of Faculty of Pharmacy, Cairo University*.
9. Beenus S C, Hosus M C, Talawas M B and Laddi U V (1994), "Synthesis and Evaluation of Antimicrobial Activity of Some New 1, 3, 4-oxadiazoles", *Indian Journal of Heterocyclic Chemistry*, Vol. 3, pp. 237-239.
10. Boström J, Hogner A, Llinàs A, Wellner E and Plowright A T (2012), "Oxadiazoles in Medicinal Chemistry", *J. Med. Chem.*, Vol. 55, pp. 1817-1830.
11. Boström J, Hogner A, Llinàs A, Wellner E and Plowright A T (2012), "Oxadiazoles in Medicinal Chemistry", *J. Med. Chem.*, Vol. 55, pp. 1817-1830.
12. Cai-Jun Chen, Bao-An Song, Song Yang, Guang-Fang Xu, Pinaki S Bhadury, Lin-Hong Jin, De-Yu Hu, Qian-Zhu Li, Fang Liu, Wei Xue, Ping Lu and Zhuo Chen (2007), *Bioorganic & Medicinal Chemistry*, Vol. 15, pp. 3981-3989.
13. Chandrakantha B, Shetty Prakash, Nambiyar Vijesh, Isloor Nishitha and Isloor Arun M (2010), "Synthesis, Characterization and Biological Activity of Some New 1, 3, 4-oxadiazole Bearing 2-Flouro-4-Methoxy Phenyl Moiety", *European Journal of Medicinal Chemistry*, Vol. 45, pp. 1206-1210.
14. Chen C, Song B, Yang S, Xu G, Bhadury P S, Jin L, Hu D, Li Q, Liu F, Xue W, Lu P and Chen Z (2007), "Synthesis and Antifungal Activities of 5- (3, 4, 5-trimethoxyphenyl)-2-sulfonyl-1, 3, 4-thiadiazole and 5-(3, 4, 5-trimethoxyphenyl)-2-sulfonyl-1, 3, 4-oxadiazole Derivatives", *Bioorg. Med. Chem.*, Vol. 15, pp. 3981-3989.
15. Chen C, Song B, Yang S, Xu G, Bhadury P S, Jin L, Hu D, Li Q, Liu F, Xue W, Lu P and Chen Z (2007), "Synthesis and Antifungal Activities of 5- (3, 4, 5-trimethoxyphenyl)-2-sulfonyl-1, 3, 4-thiadiazole and 5-(3, 4, 5-trimethoxyphenyl)-2-sulfonyl-1, 3, 4-oxadiazole Derivatives", *Bioorg Med Chem*, Vol. 15, pp. 3981-3989.
16. Demirbas A, Sahin D, Demirbas N, Karaoglu S A and Bektas H (2010), "Synthesis and Antimicrobial Activities of 2-(5-mercapto)-1, 3-oxadiazol-2-ylmethyl-1, 2, 4-triazol-3-one Derivatives", *Turk J Chem*, Vol. 34, pp. 347-358.
17. Dhansay Dewangan, Alok Pandey, Sivakumar T, Rajavel R and Ravindra Dhar Dubey (2010), "Synthesis of Some Novel 2, 5-Disubstituted 1, 3, 4-Oxadiazole and its Analgesic, Anti- Inflammatory, Anti-Bacterial and Anti-Tubercular Activity", *International Journal of ChemTech Research CODEN(USA): IJCRGG*, ISSN: 0974-4290, Vol. 2, No. 3, pp. 1397-1412.

18. Formagio A S N, Tonin L T D, Foglio M A, Madjarof C, Carvalho J E D and Costa W F A (2008), "Synthesis and Antitumoral Activity of Novel 3-(2-substituted-1, 3, 4-oxadiazol-5-yl) and 3-(5-substituted-1, 2, 4-triazol-3-yl) b-carboline Derivatives", *Bioorg Med Chem*, Vol. 16, pp. 9660- 9667.
19. Fuloria N K, Singh V and Ali M (2009), "Synthesis and Antimicrobial Evaluation of Some New Oxadiazoles Derived from Phenylpropionoylhydrazides", *Molecules*, Vol. 14, No. 5, pp. 1898-1903.
20. Fuloria N K, Singh V, Shaharyar M and Ali M (2009), "Synthesis and Antimicrobial Evaluation of Some New Oxadiazoles Derived from Phenylpropionohydrazides", *Molecules*, Vol. 14, pp. 1898-1903.
21. Hiroyuki Watanabe, Masahiro Ono, Ryoichi Ikeoka, Mamoru Haratake, Hideo Saji and Morio Nakayama (2009), "Synthesis and Biological Evaluation of Radioiodinated 2, 5-diphenyl-1, 3, 4-oxadiazoles for Detecting b-Amyloid Plaques in the Brain", *Bioorganic & Medicinal Chemistry*, Vol. 17, pp. 6402-6406.
22. Husain A, Ahuja P and Sarafroz (2009), "Synthesis and Biological Evaluation of b-aryolpropionic Acid Based 1, 3, 4-oxadiazoles", *Indian J Pharm Sci*, Vol. 71, pp. 62-79.
23. James T Palmer, Bernard L Hirschbein, Harry Cheung, John McCarter, James W Janc, Walter Yua Z and Gregg Wesolowskib (2006), "Keto-1, 3, 4-oxadiazoles as Cathepsin K Inhibitors", *Bioorganic & Medicinal Chemistry Letters*, Vol. 16, pp. 2909-2914.
24. Jayamma Y, Sarangapani M and Reddy V M (1996), "Synthesis and Evaluation of Biological Activity of Novel Heterocycles", *Indian Journal of Heterocyclic Chemistry*, Vol. 6, pp. 111-114.
25. Jayashankar B, Lokanath Rai K M, Baskaran N and Sathish H S (2009), "Synthesis and Pharmacological Evaluation of 1, 3, 4-oxadiazole Bearing Bis (Heterocycle) Derivatives as Anti-Inflammatory and Analgesic Agents", *European Journal of Medicinal Chemistry*, Vol. 44, pp. 3898-3902.
26. Kakanejadifard A, Azarbani F, Zabardasti A, Kakanejadifard S, Ghasemian M, Esnaashari F, Omid S, Shirali S and Rafieefar M (2013), "The Synthesis, Structural Characterization and Antibacterial Properties of Some 2-((4-amino-1, 2, 5-oxadiazol-3-ylimino) methyl)-4-(phenyldiazenyl) Phenol", *Dyes and Pigments*, Vol. 97, pp. 215-221.
27. Karthikeyan M S, Parsad D J, Mahalinga M, Holla B S and Kumari N S (2008), "Antimicrobial Studies of 2, 4-dichloro-5-fluorophenyl Containing Oxadiazoles", *Eur J Med Chem*, Vol. 43, No. 1, pp. 25-31.
28. Kucukguzel S G, Oruc E E and Rollas S (2002), "Synthesis, Characterization and Biological Activity of Novel 4-thiazolidinones, 1, 3, 4-oxadiazoles and Some Related Compounds", *European Journal of Medicinal Chemistry*, Vol. 37, No. 3, pp. 197-206.
29. Kumar R and Khokara S L (2012), "Chemistry and Common Synthetic Route of 1, 3, 4-oxadiazole: An Important Heterocyclic Moiety in Medicinal Chemistry", *Int J Inst Ph Life Sci*, Vol. 2, No. 5, pp. 126-135.
30. Lohman Sui Ping Lee, Asra Malikzay, James Fleming, Jason Gerlak, Ying Wang, Robin L

- Rosler, Kai Zhou, Stan Mitelman, Margarita Camara, David Surguladze, Jacqueline F Doody and Carolina Tuma M (2006), "Bioorganic & Medicinal Chemistry Letters", Vol. 16, pp. 1191-1196.
31. Mayekar A N, Yathirajan H S, Narayana B, Sarojini B K and Kumari N S (2010), "Synthesis and Antimicrobial Studies on New Substituted 1, 3, 4-Oxadiazole Derivatives Bearing 6-bromonaphthalene Moiety", *Int J Chem*, Vol. 2, No. 2, pp. 191-201.
32. Mayekar A N, Yathirajan H S, Narayana B, Sarojini B K and Kumari N S (2010), "Synthesis and Antimicrobial Studies on New Substituted 1, 3, 4-Oxadiazole Derivatives Bearing 6-Bromonaphthalene", *Moiety. Int J Chem*, Vol. 2, No. 2, pp. 191-201.
33. Mishra M K, Gupta A K, Negi S and Bhatt M (2010), *Int. J. Pharma Sciences and Reserch*, Vol. 1, No. 3, pp. 172-177.
34. Mohammad Shaharyar and Mohammad Wasim Akhter (2009), *Acta Poloniae Pharmaceutica n Drug Research*, Vol. 66, No. 4, pp. 393-397.
35. Mohammed Afroz Bakht, Shahar Yar M, Sami Gaber Abdel-Hamid, Saleh I Al Qasoumi and Abdul Samad (2010), "Molecular Properties Prediction, Synthesis and Antimicrobial Activity of Some Newer Oxadiazole Derivatives", *European Journal of Medicinal Chemistry*, Vol. 45, pp. 5862-5869.
36. Morihisa Saitoh, Jun Kunitomo, Eiji Kimura, Yoji Hayase, Hiromi Kobayashi, Noriko Uchiyama, Tomohiro Kawamoto, Toshimasa Tanaka, Clifford D Mol, Douglas R Dougan, Garret S Textor, Gyorgy P Snell and Fumio Itoh (2009), "Design, Synthesis and Structure-Activity Relationships of 1, 3, 4-oxadiazole Derivatives as Novel Inhibitors of Glycogen Synthase Kinase-3b", *Bioorganic & Medicinal Chemistry*, Vol. 17, pp. 2017-2029.
37. Morihisa Saitoh, Jun Kunitomo, Eiji Kimura, Yoji Hayase, Hiromi Kobayashi, Noriko Uchiyama, Tomohiro Kawamoto, Toshimasa Tanaka, Clifford D Mol, Douglas R Dougan, Garret S Textor, Gyorgy P Snell and Fumio Itoh (2009), "Design, Synthesis and Structure-Activity Relationships of 1, 3, 4-oxadiazole Derivatives as Novel Inhibitors of Glycogen Synthase Kinase-3b", *Bioorganic & Medicinal Chemistry*, Vol. 17, pp. 2017-2029.
38. Nesynov E P and Grekov A P (1964), "The Chemistry of 1, 3, 4-oxadiazole Derivatives", *Russ Chem Rev*, Vol. 33, No. 10, pp. 508-515.
39. Nitin Bhardwaj, Saraf S K and Pankaj Sharma (2009), "Pradeep Kumar, Syntheses", *E-Journal of Chemistry*, Vol. 6, No. 4, pp. 1133-1138.
40. Om Prakash, Manoj Kumar, Rajesh Kumar, Chetan Sharma and Aneja K R (2010), "Hypervalent Iodine(III) Mediated Synthesis of Novel Unsymmetrical 2, 5-disubstituted 1, 3, 4-oxadiazoles as Antibacterial and Antifungal Agents", *European Journal of Medicinal Chemistry*, Vol. 45, pp. 4252-4257.
41. Ouyang X, Piatnitski E L, Pattaropong V, Chen X, He H Y, Kiselyov A S, Velankar A, Kawakami J, Labelle M, Smith L, et al. (2006), "Oxadiazole Derivatives as a Novel Class of Antimitotic Agents: Synthesis, Inhibition of Tubulin Polymerization, and Activity in Tumor Cell Lines", *Bioorg. Med. Chem. Lett.*, Vol. 16, pp. 1191-1196.

42. Parkash O, Kumar M, Sharma C and Aneja K R (2010), "Hypervalent Iodine(III) Mediated Synthesis of Novel Unsymmetrical 2, 5-disubstituted 1, 3, 4-oxadiazoles as Antibacterial and Antifungal Agents", *Eur. J. Med. Chem.*, Vol. 45, No. 9, pp. 4252-4257, doi: 10.1016/j.ejmech.2010.06.023.
43. Pevear D, Tull T, Seipel M and Groarke J (1999), "Activity of Pleconaril Against Enteroviruses", *Antimicrob Agents Chemother*, Vol. 43, No. 9, pp. 2109-2015.
44. Reddy C S, Rao L S, Devi M V and Nagaraj A (2012), "Synthesis and Antimicrobial Activity of Linked Heterocyclics Containing Pyrazole and Oxadiazoles", *J Heterocyclic Chem*, Vol. 49, p. 1038.
45. Savarino A (2006), "A Historical Sketch of the Discovery and Development of HIV-1 Integrase Inhibitors", *Expert Opin Investig Drugs*, Vol. 15, No. 12, pp. 1507-1522.
46. Shaker R M, Mahmoud A F and Abdel-Latif F F (2005), "Synthesis and Biological Activities of Novel 1, 4-bridged bis-1, 2, 4-triazoles, bis-1, 3, 4-thiadiazoles and bis-1, 3, 4-oxadiazoles", *Phosphorus, Sulfur, and Silicon*, Vol. 180, pp. 397-406.
47. Shridhar A H, Keshavayya J, Peethambar S K and Hoskeri H Joy (2012), "Synthesis and Biological Activities of Bis Alkyl 1, 3, 4-oxadiazole Incorporated Azo Dye Derivatives", *Arabian Journal of Chemistry*, Vol. 014, pp. 741-758.
48. Somani R R, Agrawal A G, Kalantri P P, Gavarkar P S and Clercq E D (2011), "Investigation of 1, 3, 4-Oxadiazole Scaffold as Potentially Active Compounds", *IJDDD*, Vol. 2, No. 1, pp. 353-360.
49. Somani Rakesh R, Agrawal Anuj G, Kalantri Pushkar P, Gavarkar Pratibha S and Clercq Erik De (2011), "Investigation of 1, 3, 4-Oxadiazole Scaffold as Potentially Active Compounds", *International Journal of Drug Design and Discovery*, Vol. 2, No. 1, pp. 353-360.
50. Tully W R, Gardner C R, Gillespie R J and Westwood R (1991), "2-(oxadiazolyl)-and 2-(thiazolyl) imidazo [1, 2-a] Pyrimidines as Agonists and Inverse Agonists at Benzodiazepine Receptors", *J Med Chem*, Vol. 34, No. 7, pp. 2060-2206.
51. Usman Ghani and Nisar Ullah (2010), "New Potent Inhibitors of Tyrosinase: Novel Clues to Binding of 1, 3, 4-thiadiazole-2(3H)-thiones, 1, 3, 4-oxadiazole-2(3H)-thiones, 4-amino-1, 2, 4-triazole-5(4H)-thiones and Substituted Hydrazides to the Dicopper Active Site", *Bioorganic & Medicinal Chemistry*, Vol. 18, pp. 4042-4048.
52. Virginija Jakubkiene, Milda Malvina Burbuliene, Giedrute Mekuskiene, Emilija Udreinaite, Povilas Gaidelis and Povilas Vainilavicius (2003), "Synthesis and Anti-Inflammatory Activity of 5-(6-methyl-2-substituted 4-pyrimidinylloxymethyl)-1, 3, 4-oxadiazole-2-thiones and their 3 Morpholinomethyl Derivatives", *Il Farmaco*, Vol. 58, pp. 323-328.
53. Zheng X, Li Z, Wang Y, Chen W, Huang Q, Liu C and Song G (2003), "Synthesis and Insecticidal Activities of Novel 2, 5-disubstituted 1, 3, 4-oxadiazoles", *J. Fluorine Chem.*, Vol. 123, pp. 163-169.